

## ROLE OF ORAL PANTOPRAZOLE AS PRE OPERATIVE PREANAESTHETIC PREMEDICATION FOR THE PROPHYLAXIS OF ACID ASPIRATION IN ELECTIVE ADULT SURGERY

Altaf Hussain, Abdul Hamid Hasan Al-Saeed, Syed Shahid Habib\*, Mansoor Aqil, Sayeed Nawaz, Adeel-ur-Rahman

Department of Anaesthesiology (41), King Khalid University Hospital, \*Department of Physiology, College of Medicine & King Khalid University Hospital, Al-Riyadh, Saudi Arabia.

**Background:** The effect of drugs affecting gastric pH and volume has been studied extensively but the effect of duodenogastric reflux on gastric pH and volume at the same time has not been evaluated. **Patients and Methods:** This prospective, triple blind, randomized and placebo controlled clinical trial was conducted on 108 adult inpatients of American Society of Anaesthesiologist physical status I–II, and aged 15–70 years. The patients in Group C (control) received placebo while Group P (Pantoprazole) orally at 9:00 PM, a night before elective surgery. On the next day, gastric contents were aspirated with a large bore, multi-orifices gastric tube passed through an endotracheal tube placed blindly in oesophagus after tracheal intubation and analysed for pH, volume and the presence of bile salts. **Results:** Thirty (28.57%) samples out of 105 were contaminated with duodenal contents and 2 with blood. Duodenogastric reflux significantly affected pH and volume of gastric contents in subgroups (C-1 versus C-2: *p*-value for pH (0.0009) and volume (0.0236) and P-1 versus P-2: *p*-value for pH (0.0348) and volume (0.0003). Pantoprazole, after excluding samples contaminated with duodenogastric refluxate, increased pH (*p* 0.0118), decreased volume (*p* 0.0009) and the proportion of the patients (*p* 0.0324) considered “at risk” compared with Placebo. **Conclusion:** Preoperative oral administration of Pantoprazole 40 mg reduced residual gastric content volume  $\leq 25$  ml and increased pH  $\geq 2.5$ , possibly reducing the effects of pulmonary aspiration of gastric contents.

**Keywords:** Aspiration, duodenogastric refluxate, gastric pH, volume, Pantoprazole

### INTRODUCTION

Aspiration has been defined as ‘inhalation of material into the airway below the level of the true vocal cords’. The severity of aspiration pneumonitis depends upon pH and volume of gastric juice aspirated.<sup>1</sup> Acid-suppressing agents are used in the perioperative setting to reduce the risk of lung injury from aspiration of gastric contents. Acid-induced lung injury is associated with a high mortality—approximately 40–50%.<sup>2</sup> Although risk factors for acid aspiration continue to be debated, the literature suggests that patients with a gastric pH of  $< 2.5$  and a volume of at least 25 ml are at greatest risk.<sup>3</sup>

Pantoprazole, a proton pump inhibitor, is used in peptic ulcers and other acid dyspeptic disorders of upper gastrointestinal tract in a dose of 40 mg orally once daily.<sup>4</sup> The effect of single oral dose of Pantoprazole 40 mg on preoperative gastric fluid pH and volume has not yet been studied.

Our aim of study was whether duodenogastric reflux<sup>5</sup> [transpyloric retrograde flow of alkaline duodenal contents into the acidic gastric contents] significantly affect pH and volume of gastric contents and if yes, then whether a single oral dose of Pantoprazole 40 mg, administered a night before surgery, brings the pH  $\geq 2.5$  and volume  $\leq 0.4$  ml/Kg or  $\leq 25$  ml in adult patients undergoing elective surgery by excluding cases contaminated with duodenogastric

reflux. Such samples do not represent true gastric contents rather alkaline duodenal fluid mixed with acidic gastric contents.

### PATIENTS AND METHODS

The study protocol was approved by the University Hospital Research and Ethics Committee and written; informed consent was obtained from all the patients.

#### Patients and Group Assignment

We explored the effect of single oral dose of Pantoprazole 40 mg administered at 9:00 PM, a night before elective surgery, on intragastric pH and volume in adult 108 inpatients of American Society of Anaesthesiologists (ASA) physical status I–II, aged 15–70 years to be intubated with cuffed endotracheal tube.

Obese patients of Body Mass Index (BMI) more than 40 Kg/m<sup>2</sup>, patients with upper gastrointestinal disorders, receiving medications known to affect the composition and volume of the stomach contents, Mallampati class IV and/or mouth opening less than 5 Cm and/or thyromental distance less than 6.5 Cm and/or history of documented difficult intubation, Diabetes Mellitus, intestinal obstruction and parturients were excluded from the study. Patients whose gastric aspirates contained bile salts due to duodenogastric reflux (DGR) or gastric contents were mixed with blood in the gastric tube were not included in the final statistical analysis while analysing pH and volume of

gastric contents because these samples are not true gastric contents rather alkaline duodenal fluid mixed with acidic gastric contents or blood mixed with gastric contents.

Patients were randomly assigned to one of the two groups by sealed enveloped method as follows (n=54 in each group): Group C (Placebo) and Group P (Pantoprazole). Age, sex, weight, height, BMI, ASA physical status, and the drug given were recorded for each patient. The medications were given orally with 20 ml of drinking water at 9:00 PM, a night before elective surgery. All patients also took oral diazepam 10 mg at the same time. All patients, according to the Hospital policy, remained nil per os (NPO) after 12 AM. Upon arrival in the waiting area of the operating room, patients were asked if they had been aware of any unusual feelings (side effects) like headache, gastrointestinal upset etc. after taking the medications, a night before elective surgery. This was also noted.

#### Collection and Analysis of Gastric Contents

After pre-oxygenation with 100% O<sub>2</sub> by face mask using four breaths vital capacity method, general anaesthesia was induced with injection fentanyl 1–2 µg/Kg, propofol 2–3 mg/Kg and rocuronium 0.6–0.9 mg/Kg. The lungs were ventilated taking care not to inflate the stomach. Maintaining cricoid pressure, trachea was intubated with cuffed endotracheal tube. All inductions were uneventful. After tracheal intubation, an endotracheal tube sized 8.0 mm internal diameter coated with paraffin liquid internally was passed orally in the oesophagus with anterior displacement of larynx. A predetermined length marked with adhesive tape (from Xiphoid process to ear lobules- and from ear lobules to nasal tip) of stomach tube<sup>6</sup> (Jamjoom Medical Industries, Jeddah, Saudi Arabia) sized 16 F was passed through this oesophageally placed endotracheal tube<sup>7</sup>. Placement of orogastric tube within the stomach was confirmed by auscultation over the epigastrium by injecting 10–15 ml of air. Gastric contents were gently aspirated manually with 60 ml of syringe by an investigator who was unaware of the group assignment. Applying and maintaining manual pressure over the epigastrium while the patient was in supine and then left and right lateral positions, orogastric tube was then manipulated to ensure maximum emptying of gastric contents. The orogastric tube was taken out followed by endotracheal tube placed in oesophagus. Any problem encountered during inserting or removing either the oesophageally placed endotracheal tube or orogastric tube was also noted. The gastric contents were visually inspected for blood. The volume of gastric contents was measured with graduated syringe and pH with pH meter (Model 215 version 3.4, Denver Instrument Company, United States). The pH meter was calibrated using standard buffers at pH values of 4, 7 and 9.20. This pH meter has a precision of 0.01 units over the entire pH

range. A minimum of 1 ml. volume of gastric contents was sufficient for pH determination with pH meter. Samples less than 1 ml. were considered as no gastric contents because a minimum volume of 1 ml. of gastric contents was sufficient for pH-meter. Using bile salts as a marker for bile, we applied qualitative Hay's Sulphur test to detect bile salts. A minimum volume of 1 ml. of gastric contents was sufficient to perform Hay's Sulphur test. In this test finely powered Sulphur is sprinkled upon the surface of cool (17 °C or below) liquid. If bile salts are present Sulphur sinks down, sooner or later, in accordance with their percentage. If bile salts are present in from 1:5000 (0.02% or 200 µg/ml) to 1:10,000 (0.01% or 100 µg/ml) Sulphur at once begins to sink and all precipitated in 2 or 3 minutes; even in a dilution of 1:120,000 (0.0008% or 8.33 µg/ml) precipitation occurs. On the other hand if Sulphur remains floating on the surface, bile salts are absent.<sup>8</sup>

Time since premedication, time since *nil per os* (NPO), pH, volume of gastric contents and result of Hay's Sulphur test were also recorded for each patient. On the basis of Hay's Sulphur test, we further subdivided the Group C into Subgroups C-1 and C-2 and Group P into Subgroups P-1 and P-2 to observe the effect of duodenogastric refluxate on pH and volume of gastric aspirates.

#### Statistical Analysis

Statistical tests were performed using GraphPad Software, Inc., San Diego, United States, and results are expressed as absolute values (percentage) or Mean±SD.

Statistical comparisons between the two Groups C and P were carried out using two-tailed Student's (unpaired) t test for age, weight, height, BMI, time since premedication, time since NPO, pH and volume (Subgroups C-1, C-2, P-1 and P-2).

Fisher's exact test (two-tailed) was applied for sex, ASA physical status and risk of aspiration according to the criteria defined by Roberts and Shirley<sup>9</sup> (pH ≤2.5 and volume ≥0.4 ml/Kg or 25 ml). A p- value of less than 0.05 deemed statistically significant.

Power analysis revealed that the sample size (n= 30 in each group) of the study is sufficient to detect medium differences ([mean 1-mean 2]/SD= 0.5–0.7) in variables (pH and volume) at a significance level of 0.05 with the power of 0.6–0.8.<sup>10</sup>

## RESULTS

One hundred and eight (108) adult inpatients undergoing elective General (n=54), Orthopaedic (n=25), Gynaecological (n=11), Urology (n=9), Thoracic (n=8) and Neuro (n=1) Surgery were studied. Physical characteristics of patients and timings of events are shown in Table-1. There was no statistically significant difference between the Groups C and P.

**Table-1: Clinical characteristics of patients and timings of events.**

Values are expressed as Mean±SD or numbers (percentage).

Physical characteristics of patients	Group C n=54	Group P n=54	p-Value
Age (years)	33.57 ±12.45	34.65 ±12.97	0.6616
Sex			
Male	27 (50%)	28 (51.85%)	1.000
Female	27 (50%)	26 (48.14%)	
ASA Physical Status			
Class-I	43 (79.62%)	38 (70.37%)	0.3743
Class-II	11 (20.38%)	16 (29.62%)	
Weight (Kg)	74.59 ±14.29	75.99 ±17.47	0.6482
Height (Cm)	162.10 ±7.79	163.25 ±10.14	0.5132
Body Mass Index (Kg/m <sup>2</sup> )	28.52 ±5.57	28.44 ±5.43	0.9442
Timings of events			
Time since premedication (Min)	816.33 ±118.13	835.15 ±107.27	0.3882
Time since NPO (Min)	643.11 ±127.02	651.44 ±101.46	0.7072

We obtained gastric contents of 107 patients. One patient in Group P has no gastric contents while two samples (one from each Group) were mixed with blood. Hay's test was performed on 105 samples and was positive in 30 (28.57%) patients, 16 (6 males and 10 females) in Group C and 14 (7 males and 7 females) in Group P. The average (range) pH and volume of contaminated cases with duodenal contents were 5.75 (1.63–6.98) and 39.86 (5.0–96.0) ml and with blood were 7.13 (6.53–7.73) and 3.5 (3.0–4.0) ml. Samples contaminated with either duodenal fluid (30) or blood (2) were not considered as true gastric contents and not included in the final statistical analysis while analyzing pH and volume of gastric contents.

The pH and volume of all the Subgroups is shown in Table-2. Duodenogastric refluxate significantly affected both the pH and volume of gastric contents in both the Groups C (*p*-values 0.0009 and 0.0236, respectively) and Group P (*p*-value 0.0348 and 0.0003, respectively). The pH and volume of true gastric contents (Subgroups C-2 and P-2 is also shown in Table-2. Pantoprazole significantly affected both the pH (*p*=0.0118) and volume (*p*=0.0009) of gastric contents compared with placebo Group.

The proportion of the patients considered 'at risk' of significant lung injury should aspiration occur is shown in the Table-3 (after excluding contaminated samples with duodenogastric refluxate). There was a statistically significant difference between the Groups C-2 and P-2 (*p* 0.0324). No side effect of study drug

was noted. All patients were discharged from the hospital without any problem.

**Table-2: pH and volume of gastric contents (Mean±SD)**

Variables	Group C n=54		Group P n=54	
	Group C-1 n=53	Group C-2 n=37	Group P-1 n=52	Group P-2 n=38
pH	2.85±1.90	1.75±0.47	3.73±2.09	2.85±1.69
Volume (ml)	25.79±18.61	18.09±9.68	19.32±14.34	12.68±8.43

Note: Samples contaminated either with blood (2) or no gastric contents (1) are not included in Subgroups C1 and P1. Group C-1 and Group P-1 include contaminated samples with duodenogastric refluxate as well. Group C-2 and Group P-2 represent true gastric contents. Comparison between the Subgroups *p*-value for pH (0.0009) and volume (0.0236) between Group C-1 and Group C-2. *p*-value for pH (0.0348) and volume (0.0003) between Group P-1 and Group P-2. *p*-value for pH (0.0118) and volume (0.0009) between Group C-2 and Group P-2.

**Table-3: Patients at risk according to defined criteria (n (%))**

Variables	Group C-2 n=37	Group P-2 n=38	p-value
Patients with pH ≤2.5	35 (94.59%)	23 (60.52%)	0.0006
Patients with volume ≥25 ml.	13 (35.13%)	8 (21.05%)	0.2018
Patients with pH ≤2.5 and volume ≥25 ml.	13 (35.13%)	5 (13.15%)	0.0324

Note: Samples mixed either with duodenal contents (30) or blood (2) or having no contents (1) are not included in Subgroups C-2 and P-2.

## DISCUSSION

Many pharmacological attempts, including the use of H<sub>2</sub>-receptor antagonists, proton pump inhibitors (PPIs) and antacids have been made to reduce the risk of pulmonary aspiration by decreasing acidity and volume of gastric fluid.<sup>9</sup> Pantoprazole, developed in Germany in 2000, is the third proton pump inhibitors after omeprazole and Lansoprazole<sup>11</sup> used in clinical practice. Proton pump inhibitors act by irreversibly blocking the hydrogen/potassium adenosine triphosphatase enzyme system (the H<sup>+</sup>/K<sup>+</sup> ATPase, or more commonly just gastric proton pump) of the gastric parietal cell. The proton pump is the terminal stage in gastric acid secretion, being directly responsible for secreting H<sup>+</sup> ions into the gastric lumen, making it an ideal target for inhibiting acid secretion.

There is only one study available that was conducted to evaluate the effect of intravenous Pantoprazole and ranitidine for improving preoperative gastric fluid properties in adults undergoing elective surgery.<sup>12</sup> In this study all drugs were given intravenously one hour before surgery. In our study the Mean±SD, pH, volume and proportion of patients at risk are 2.85±1.69, 12.68±8.43 ml and 13.15% compared with intravenous Pantoprazole 5.30±1.84 (*p*<0.0001), 15.20±15.52 ml (*p*=0.7166) and 10% (*p*=1.0000), respectively. The primary outcome of the study, the proportion of patients at risk is almost the same. A better aspect in our study may be the exclusion of all

duodenogastric refluxate contaminated samples. In this current study 30 samples out of 105 were positive for bile salts in their gastric aspirates. This is, no doubt, a big number and if unfortunately, patient aspirates it will damage the lungs as well. We should have included a prokinetic agent to get rid of duodenogastric refluxate. More work is required in this regard.

Mendelson CL described the aspiration of gastric contents (Mendelson's syndrome) in 1946 in obstetrical cases.<sup>13</sup> Since then extensive work has been done and reported in anaesthesia literature. In all the previous studies conducted, importance of duodenogastric reflux (DGR), as a possible factor that can affect both the pH and volume of gastric contents, has never been addressed. Duodenal contents consist of bile, pancreatic juice, small intestine and Brunner's gland secretion. All these secretions are alkaline in nature due to  $\text{HCO}_3^-$  ions.<sup>14</sup> When duodenal contents flow in retrograde fashion, then mix with acid and Pepsin<sup>5</sup> in the stomach and bring the pH towards less acidity thus affecting pH and at the same time increase the volume of gastric contents similar to oral ingestion of sodium citrate. To overcome this problem, first, we aspirated gastric contents in optimal position of the patient as described by Niinai *et al.*<sup>15</sup> Secondly, we passed a predetermined length of stomach tube so that it should not go beyond pyloric sphincter. Thirdly, we excluded those samples that were positive for Hay's Sulphur test while analysing final pH and volume of gastric contents.

In this current study, we passed gastric tube through an endotracheal tube passed blindly in the oesophagus. Although, this technique of passing stomach tube is old<sup>7</sup>, but no body has utilised it for sampling gastric contents in any previous study. Importantly this technique can avoid contamination of gastric contents with pooled saliva in pharynx during inserting, manipulating or removing gastric contents.<sup>16</sup> Insertion of oropharyngeal airway, act of laryngoscopy and tracheal tube insertion are the stimulants that increase the production rate of saliva and saliva pools due to the lack of swallowing reflex in pharynx.

In this study two samples were found to be mixed with blood due to gastric mucosal entrapment. Gastric mucosal entrapment occurs particularly when air and fluid has been aspirated and stomach is collapsed. Gastric mucosa is caught into the side holes of stomach even with gentle suction effect. Bleeding may occur and can be seen in stomach tube thus giving pH of blood mixed with gastric contents rather than pure gastric contents. It is commonly believed that the sump tubes (double-lumen) are more effective than the single lumen variety, but there is no scientific evidence to support this view.<sup>17</sup> However, any sample containing any amount of visible blood mixed with gastric contents was not

considered true sample and was not included in the analysis for pH, volume and Hay's Sulphur test.

We applied Hay's Sulphur test to detect bile salts in the gastric contents. This simple, sensitive and fairly reliable test depends on the principal that bile salts have the property of reducing the surface tension of fluids in which they are contained<sup>18</sup>, was devised in 1886 by Matthew Hay (1855–1932).

In this present study, total gastric volume may have been underestimated by the blind aspiration via gastric tube in each patient due to the functional divisions of the stomach into antral and fundal sacs.<sup>19</sup> A similar error would occur in all patients of both groups and inter-group comparisons are, therefore, valid. Due to difficulty in accurate measurement of gastric fluid volume, pH is a better marker for aspiration assessment.

## CONCLUSION

Oral Pantoprazole 40 mg administered a night before elective surgery does improve the gastric environment, excluding those samples contaminated with duodenogastric reflux, and at the time of induction of anaesthesia may prove to be useful in decreasing risk of aspiration due to gastric contents.

## ACKNOWLEDGEMENTS

Authors are very thankful to the laboratory staff of Clinical Chemistry Unit, Nursing Department and Colleagues of Anaesthesia and Surgical Departments of King Khalid University Hospital for their co-operation.

## REFERENCES

1. Mikawa K, Nishina K. Preoperative medication. In: Miller RD, ed. Atlas of anesthesia. Vol 3. Churchill Livingstone; Philadelphia, 1997.p1–24.
2. Matthey MA, Rosen GD: Acid aspiration induced lung injury: New insights and therapeutic options (Editorial). Am J Respir Crit Care Med 1996;154:277–8.
3. Stuart JC, Kan AF, Rowbottom SJ, Yau G, Gin T: Acid aspiration prophylaxis for emergency caesarean section. Anaesthesia 1996;51:415–21.
4. Christopher Haslet, Edwin R. Chilvers, John A. A. Hunter and Nicholas A. Boon. Davidson's Principles and Practice of Medicine. 18<sup>th</sup> Ed. Churchill Livingstone; Edinburgh, 1999.p635.
5. Joel E, Richter. Duodenogastric reflux –induced (alkaline) esophagitis. Curr Treat Options Gastroenterol 2004;7:53–8.
6. McConnell EA. Ten problems with nasogastric tubes and how to solve them. Nursing 1979;9:78–81.
7. Siegel IB, Kahn RC. Insertion of difficult nasogastric tube through a naso-esophageally endotracheal tube. Crit Care Med 1987;15:876–7.
8. John Dixon Mann. Physiology and Pathology of Urine. Griffin.1913;227.
9. Roberts RB, Shirley MA. Reducing the risk of acid aspiration during cesarean section. Anesth Analg 1974;53:859–68.
10. Kahoru N, Katsuya M, Nobuhiro M, Yumiko T, Makato S, Hidefumi O. A comparison of Lansoprazole, omeprazole and ranitidine for reducing preoperative gastric secretion in adult patients undergoing elective surgery. Anesth Analg 1996;82(4):832–6.

11. Bardhan KD. Pantoprazole: A new proton pump inhibitors in the management of upper gastrointestinal disease. *Drugs Today* 1999;35:773–808.
12. Dilek M, Alparslan T, Beyhan K, Pinar S, Mevlut T, Zafar P. The effect of intravenous Pantoprazole and ranitidine for improving preoperative gastric fluid properties in adults undergoing elective surgery. *Anesth Analg* 2003;97:1360–3.
13. Mendelson CL. The aspiration of stomach contents into the lungs during obstetric anesthesia. *Am J Obst and Gynecol* 1946;52:191–205.
14. Guyton AC. *Textbook of Medical Physiology*. 10<sup>th</sup> Ed. W.B. Saunders, Inc.; Philadelphia, 2000. p738–53.
15. Niinai H, Nakao M, Nakatani K, Kawaguchi R, Takezaki T, Kobayashi N. Significance of patient's position in measuring gastric contents. *Masui* 1994;43(11):1665–7.
16. Altaf H, Al-Saeed AH, Habib SS. Effect of Saliva on pH and volume of gastric contents while sampling from stomach with two different techniques of orogastric intubation. *The Internet Journal of Anesthesiology* 2007. URL: <http://www.britannica.com/bps/additionalcontent/18/25939928/Effect-of-Saliva-on-pH-and-volume-of-gastric-contents-while-sampling-from-stomach-with-two-different-techniques-of-orogastric-intubation>
17. Ikard RW, Federspiel CE. A comparison of Levi and sum nasogastric tubes for postoperative gastrointestinal decompression. *Am J Surg* 1987;53:50–3.
18. Joshi A, Rashmi. *A Textbook of Practical Biochemistry*. B. Jain Publishers; New Delhi, 2004. p51.
19. Holdsworth JD, Johnson K, Mascall G, Roulston RG, Tomlinson PA. Mixing of antacids with stomach contents. Another approach to the prevention of the acid aspiration (Mendelson's Syndrome). *Anaesthesia* 1980;35:461–50.

---

### Address for Correspondence:

**Dr. Altaf Hussain**, Consultant Anaesthetist, Department of Anaesthesiology (41), Post Box No. 7805, King Khalid University Hospital, Al-Riyadh 11472, Saudi Arabia. Tel: +966-1-4671597, 4671599, Fax: +966-1-4679364  
**E-mail:** draltaf94@yahoo.com